

Life history traits and sensitivity to landscape change: the case of birds and mammals of mediterranean Chile

Rasgos de historia de vida y sensibilidad a los cambios de paisaje:
el caso de los mamíferos y aves de Chile mediterráneo

RODRIGO A. VASQUEZ¹ and JAVIER A. SIMONETTI²

Departamento de Ciencias Ecológicas, Facultad de Ciencias, Universidad de Chile, Casilla 653,
Santiago, Chile, ¹e-mail: rvasquez@abello.dic.uchile.cl, ²e-mail: jsimonet@abello.dic.uchile.cl

ABSTRACT

Life history traits may constitute adequate indicators of the sensitivity of a species to changes in habitat/landscape availability. In this study we examine the role of life history traits in the responses of birds and mammals from the mediterranean-climate region of Chile to landscape change. Through a literature survey, we assess an index of sensitivity that may unravel which life history traits induce higher vulnerability to habitat/landscape transformation. Results revealed a broad range of relevant life history traits in mediterranean rodents and birds. Species with high sensitivity indices include *Chelemys megalonyx*, *Octodon bridgesi* and *O. lunatus* among rodents, and *Columba araucana*, *Patagona gigas*, *Campephilus magellanicus*, *Scelorchilus albicollis*, *Scytalopus magellanicus*, *Agelaius thilius*, *Asio flammeus*, and *Strix rufipes*, among birds. Two traits, reproductive effort and habitat requirements, are particularly important for the most sensitive species. The correlation between the sensitivity index and the conservation status of species suggests that the subjective judgments upon which Red Lists are based are supported by biological attributes. However, such listings may overlook some species. The approach reveals that a minimum of biological information can provide useful guidelines to establish criteria for the conservation of Chilean mediterranean mammals and birds.

Key words: birds, central Chile, reproductive effort, habitat, life histories, rodents.

RESUMEN

Los rasgos de historia de vida podrían constituir indicadores adecuados de la sensibilidad de las especies a los cambios en disponibilidad de hábitat/paisaje. En este estudio examinamos el papel de los atributos de historia de vida en la respuesta a los cambios de paisaje de aves y mamíferos de la región mediterránea de Chile. Para ello evaluamos un índice de sensibilidad para determinar qué rasgos de historia de vida producen una mayor vulnerabilidad a las transformaciones del hábitat/paisaje. Los resultados mostraron un amplio rango de atributos de historia de vida relevantes en roedores y aves de la zona mediterránea. Entre las especies con altos índices de sensibilidad se encontraron los roedores *Chelemys megalonyx*, *Octodon bridgesi* y *O. lunatus*, y las aves *Columba araucana*, *Patagona gigas*, *Campephilus magellanicus*, *Scelorchilus albicollis*, *Scytalopus magellanicus*, *Agelaius thilius*, *Asio flammeus*, y *Strix rufipes*. Esfuerzo reproductivo y requerimientos de hábitat son los atributos más importantes en las especies más sensibles. La correlación entre el índice de sensibilidad y el status de conservación de las especies sugiere que los juicios subjetivos sobre los cuales se basan los Libros Rojos se apoyan en atributos biológicos. Sin embargo, tales listados pueden pasar por alto algunas especies. La aproximación descrita muestra que un mínimo de información biológica puede proveer de útiles directrices para establecer criterios en la conservación de mamíferos y aves de Chile mediterráneo.

Palabras clave: aves, Chile central, esfuerzo reproductivo, hábitat, historias de vida, roedores.

INTRODUCTION

Landscape modification, mainly caused by humans, is the most common threat to species survival. Habitat loss, fragmentation and degradation usually trigger population declines, increasing the probability of a species' local and global extinction (Groom & Schumaker 1993, Fahrig & Grez 1996). Anthropogenic land changes account for 35% of contemporary animal extinctions for which causes are known. Similarly, 76% of threatened mammals are menaced by habitat destruction (Groombridge 1992). Land-use patterns are likely to intensify in the next decades (Leemans & Zuidema 1995). Consequently, criteria and indices to predict species sensitivity to landscape changes are badly needed to predict which species might be more prone to extinction.

From all phenotypic features of an organism, life history traits, being closely linked to fitness (Mousseau & Roff 1987, Roff 1997), are particularly sensitive to selective pressures, and hence to ecological changes (Stearns 1992). Life history traits have low heritability compared to other phenotypic traits (Roff 1992, 1997, Stearns 1992), and therefore may respond more rapidly to selection (Charlesworth 1994, Falconer & Mackay 1996, Roff 1997, but see Price & Schluter 1991). Ecological changes derived from landscape modifications might favour different life histories through their differential effects on reproduction, and mortality (Southwood 1977, 1988, Stearns 1992). Therefore, an analysis of life history traits, such as reproductive effort and habitat requirements, may suggest what set of attributes render a species more vulnerable to landscape change (Hansen & Urban 1992). This approach offers an alternative method to understand and predict the conservation status of species compared to the currently subjective judgments upon which most Red Lists are based (Mace 1995). By landscape change we consider habitat loss or fragmentation caused by human activities, and with an overall reduction in the amount and quality of habitat (Forman 1995).

Mediterranean evergreen shrublands are among the most disturbed and threatened biomes worldwide (Hannah et al. 1995). Mediterranean regions harbor a remarkably high biodiversity (Cowling et al. 1996); they are considered "hot-spots" precisely due to their high and unique biological richness and their intense land use, demanding urgent protection (Myers 1990). The mediterranean-climate region of Chile is no exception. It has been subjected to anthropogenic landscape changes for millennia (Fuentes 1990). This region supports a rich and diverse biota, a fraction of which either has locally vanished or face extirpation (Miller 1980, Simonetti & Cornejo 1990, Arroyo et al. 1995, Simonetti 1999, Cofré & Marquet 1999). Several mediterranean species are included in the Chilean Red List, a recognition of their threatened status (Glade 1993). However, there are no ecological comparative analyses of the biological attributes that could be underpinning their status (but see Jaksic & Jiménez 1986 for Chilean raptors).

In this paper we examine the role of life history traits in the responses of birds and mammals from the mediterranean-climate region of Chile to landscape change. Following the sensitivity index proposed by Hansen & Urban (1992) for North American bird assemblages, we aim to identify specific traits or suites of traits that contribute more significantly to species endangerment when facing landscape modifications. Second, we compare the ranking of species' sensitivity to their current conservation status in the Chilean Red List. Congruency between the sensitivity index and their conservation status will validate such listings.

MATERIAL AND METHODS

Following Hansen & Urban (1992), we use a comprehensive definition of life history, including traits related to reproductive strategy, space utilization, and habitat requirements (see also O'Connor 1985, Sibly & Calow 1985, Roff 1992). We assessed the potential responsiveness of

species to landscape modification through a sensitivity index (SI) based on six life history traits: reproductive effort (litter size times the annual number of reproductive events), type of nest or den, nest/den position, space use or abundance, habitat requirements, and migratory behavior (Table 1). These traits comprehend a range of life history attributes that include several aspects of the organismal biology of the animals, from reproductive success to resource utilization. By type of nest/den we refer to whether animals use holes and/or covered cavities in trees or soil (low vulnerability), use cup like semi-open nests/dens (intermediate vulnerability), or use open barely hollow nest/den (high vulnerability). Nest/den position represents the height from the ground (>3 m, low vulnerability; 1-3 m, intermediate; < 1 m high). In relation to migratory behavior, it has been suggested that those species most vulnerable to landscape modifications are those animals with reproductive migrations, since the lack of a habitat may generate a failed migration with no reproduction, particularly since migration interacts with other life history traits (O'Connor 1985, Hansen & Urban 1992, Sutherland & Dolman 1994). Resident non-migrant species would be least sensitive to land transformations, whereas animals with short non-

reproductive migrations (e.g., long-term traplining birds) would have an intermediate sensitive to landscape change (O'Connor 1985, Hansen & Urban 1992). Each trait could range from 1 (least sensitive) to 3 (most sensitive) depending on its value or status (Hansen & Urban 1992; Table 1). The SI for a species is obtained by summing the scores across traits. The higher the value, the more sensitive the species is to landscape change. This approach assumes that life history traits have little or no variance. While this assumption may not be entirely correct (McNamara & Houston 1996, Roff 1997), life history traits are more variable between than within species, supporting interspecific comparisons (Roff 1992, Stearns 1992).

Rather than embracing an exhaustive review of the fauna from the mediterranean region of Chile, we aim to exemplify the heuristic value of this approach focusing on a suite of species. We selected 21 bird and 11 mammal species belonging to 9 different families (6 avian and 3 mammalian), largely based on information availability. From this selected families we assessed those species that inhabit the mediterranean-climate region of central Chile, including some species that at present time

TABLE 1

Life history criteria used to rate the sensitivity (1 = least sensitive; 3 = most sensitive) of bird and mammal species to landscape change. The overall index value for a particular species is the summation of individual values across life history traits (see Material and Methods)

Crterios de historia de vida usados para categorizar la sensibilidad (1 = menos sensible; 3 = más sensible) de especies de aves y roedores a los cambios de paisaje. El valor global del índice para una especie particular es la suma de los valores individuales de cada rasgo de historia de vida (véase Material y Métodos)

Life history traits	Sensitivity score		
	1	2	3
reproductive effort (litter size/year)	> 10	6 - 10	0 - 5
nest/den form	hole	semi-open	open
nest/den height (m)	> 3	1 - 3	< 1
space use or abundance	abundant	intermediate	rare
habitat requirements	generalist	closed-canopy, open-canopy	old growth
migratory behavior	resident	short	long or reproductive

TABLE 2

Sensitivity to landscape change of (a) rodent and (b) bird species of mediterranean Chile. Higher values of the index indicate greater sensitivity. Those species with conservation status in central Chile (according to Glade 1993) are also indicated (- = without status; K = insufficiently known; I = indeterminate; V = vulnerable; E = endangered)

Sensibilidad a los cambios de paisaje de (a) roedores, y (b) aves de Chile mediterráneo. Valores mayores del índice indican mayor sensibilidad. Se indican también aquellas especies con alguna categoría de conservación para Chile central (según Glade 1993; - = no categorizada; K = insuficientemente conocida; I = indeterminada; V = vulnerable; E = en peligro)

Taxa	Family	Species	Sensitivity index	Conservation status
(a) Rodents	Muridae	<i>Abrothrix longipilis</i>	12	K
		<i>Abrothrix olivaceus</i>	10	-
		<i>Chelemys megalonyx</i>	14	E
		<i>Euneomys chinchilloides</i>	13	K
		<i>Oligoryzomys longicaudatus</i>	9	-
		<i>Phyllotis darwini</i>	11	-
	Octodontidae	<i>Octodon degus</i>	10	-
		<i>Octodon bridgesi</i>	14	V
		<i>Octodon lunatus</i>	14	V
		<i>Spalacopus cyanus</i>	12	-
Abrocomidae	<i>Abrocoma bennetti</i>	11	I	
(b) Birds	Columbidae	<i>Columba araucana</i>	15	E
		<i>Zenaida auriculata</i>	11	-
		<i>Columbina picui</i>	12	-
	Trochilidae	<i>Patagona gigas</i>	14	-
		<i>Sephanoides sephaniodes</i>	12	-
	Picidae	<i>Colaptes pitius</i>	11	-
		<i>Picoides lignarius</i>	11	-
		<i>Campephilus magellanicus</i>	14	E
	Rhinocryptidae	<i>Pteroptochos castaneus</i>	13	-
		<i>Pteroptochos megapodius</i>	13	-
		<i>Scelorchilus albicollis</i>	14	-
		<i>Scytalopus magellanicus</i>	14	-
	Icterinae	<i>Molothrus bonariensis</i>	11	-
		<i>Curaeus curaeus</i>	13	-
		<i>Agelaius thilius</i>	14	-
		<i>Sturnella loyca</i>	13	-
		Strigidae	<i>Bubo virginianus</i>	11
	<i>Glaucidium nanum</i>		10	-
<i>Asio flammeus</i>	14		K	
<i>Athene cunicularia</i>	11		-	
<i>Strix rufipes</i>	14		K	

occupy marginal areas of this region (e.g., *Campephilus magellanicus*). Our selection of families included species with disparate amounts of available information and conservation status. Therefore, the sample should be representative of any broader taxonomic analysis. Life history information was obtained through a perusal of the literature. Most of the data were obtained from Johnson (1965, 1967) for birds, and from Redford & Eisenberg (1992), and Mann (1978) for mammals, supplemented by works by Glanz (1977), Meserve & Le Boulengé (1987), and Meserve et al. (1996) for mammals, and Schlatter (1979), Jaksic & Jiménez (1986), Estades (1995), and Martínez & Jaksic (1996) for birds. Although the quality of this information is somewhat variable (owing to different sample sizes, recording methodologies, and observers, among other factors), to our knowledge, the data is the best available and sufficiently reliable to exemplify the approach. In the few cases when available information was ambiguous or scarce for a given trait, we used the value of the closest relative species with the lowest index value for that trait. Therefore, if anything, our analysis may underestimate the sensitivity to landscape change.

The outcome of the SI was compared to the ranking of each species in the Red List of Chilean vertebrates (Glade 1993). A non-parametric correlation between the index and conservation ranking was performed. The conservation status of each species in central Chile was ranked as follows: without status (i.e., unthreatened species not included in the Red List) = 1; out of danger = 2; status not defined, insufficiently known, and indeterminate = 3; rare = 4; vulnerable = 5; endangered = 6; and extinct = 7.

RESULTS

Mediterranean birds and mammals of Chile exhibit an extensive range of life history traits, with sensitivity index values from SI = 9 (of a possible minimum of 6) in the long-tailed rice rat (*Oligoryzomys*

longicaudatus Bennett, 1832) to SI = 15 (of a possible maximum of 18) in the Chilean pigeon (*Columba araucana* Lesson, 1827; Table 2). Among rodents, the murid *Chelemys megalonyx* (Waterhouse, 1845) and the octodontids *Octodon bridgesi* (Waterhouse, 1845) and *O. lunatus* (Osgood, 1943) show the highest sensitivity (SI = 14), while the long-tailed rice rat (*O. longicaudatus*), the olivaceous field mouse (*Abrothrix olivaceus* Waterhouse, 1837), the leaf-eared mouse (*Phyllotis darwini* Waterhouse, 1837), and the degu (*Octodon degus* Molina, 1782) are the least sensitive to landscape change (Table 2a).

Among birds, the Chilean pigeon (*C. araucana*) appears as the most sensitive species (SI = 15) closely followed by the Giant hummingbird (*Patagona gigas* Vieillot, 1824), the Magellanic woodpecker




Fig. 1. Correlation between sensitivity index and conservation ranking for (a) rodents, and (b) birds. Each point represents one species. Statistical analyses in the text.

Correlación entre índice de sensibilidad y nivel de conservación para (a) roedores y (b) aves. Cada punto representa una especie. Análisis estadístico en el texto.

(*Campephilus magellanicus* King, 1828), the White-throated tapaculo (*Scelorchilus albicollis* Kittlitz, 1830), the Andean tapaculo (*Scytalopus magellanicus* Gmelin, 1789), the Yellow-winged blackbird (*Agelaius thilius* Molina, 1782), and two strigiforms, the Short-eared owl (*Asio flammeus* Pontoppidan, 1763), and the Rufous-legged owl (*Strix rufipes* King, 1828), all with SI = 14 (Table 2b). Avian species with low sensitivity to landscape change comprise the Eared dove (*Zenaida auriculata* des Murs, 1847), the Chilean flicker (*Colaptes pitius* Molina, 1782), the Striped woodpecker (*Picoides lignarius* Molina, 1782), the Shiny cowbird (*Molothrus bonariensis* Gmelin, 1789), the Great horned owl (*Bubo virginianus* Gmelin, 1788), the Austral pygmy owl (*Glaucidium nanum* King, 1828), and the Burrowing owl (*Athene cucularia* Molina, 1782) (Table 2b).

In general, for the species most sensitive to habitat modification, two life history traits were particularly important: reproductive effort, and habitat requirements. Among the 11 species with $SI \geq 14$, 73% have maximum scores in reproductive effort and habitat requirements, while among the four species with $SI \leq 10$, only one species has maximum score in reproductive effort and none in habitat requirements.

The SI of both birds and mammals were significantly correlated with their rankings in the Chilean Red List (Spearman corrected for ties, $r_s = 0.86$, $P = 0.006$ for rodents; $r_s = 0.59$, $P = 0.009$, for birds; see Fig. 1), although there was a large deviance in the bird data, in particular for the lowest conservation rank (Fig. 1b).

DISCUSSION

Habitat specialization and low reproductive effort increase extinction probability (Gilpin & Soulé 1986). In fact, those life history traits were the most significant in determining the sensitivity of central Chilean species to landscape change. While this result is not surprising, its covariation and the comparatively minor effect of the

other traits, offer a guideline when selecting species for conservation efforts.

The four most sensitive species to landscape modifications require old growth forest or dense vegetation, the habitat which has been disappearing from central Chile. Among them, a paradoxical case is *O. bridgesi*, listed as Vulnerable in the Red List, but considered a pest in some commercial forests (Rodríguez 1993). Inhabitant of dense vegetation, *O. bridgesi* might use the dense understory of young plantations as an alternative habitat that could compensate the loss of natural habitat. However, this species is being controlled by poisoning, habitat modification, and predator manipulations (Murúa & Rodríguez 1989, Muñoz & Murúa 1990). The reliance of *O. bridgesi* on disappearing dense woodlands is implied by its extinction from several localities in central Chile since precolumbian times (Simonetti & Saavedra 1998).

The sensitivity of the Chilean pigeon (*C. araucana*) is primarily due also to habitat requirements (mainly old growth), low reproductive success (1-2 broods per year), and its vulnerable open nesting. Johnson (1967) stated that the Chilean pigeon nests "are mere rudimentary platforms of twigs with the light showing through in all directions". Similarly, other avian species sensitive to landscape changes, such as rhinocryptids, also require old growth forests and have low reproductive efforts. It is not surprising then, that the Magellanic woodpecker (*C. magellanicus*) shows a high SI, since it inhabits old growth forests and currently its distribution is limited to the southern boundary of the mediterranean-climate region (Araya & Millie 1988). Restricting the analysis for the same life history traits, the Rufous-legged owl (*S. rufipes*) shows the same SI as its North American relative, the Spotted owl (*S. occidentalis*; see Hanson & Urban 1992); both species are well known for their old growth forest requirements (Martínez & Jaksic 1996).

Three bird species, the Giant hummingbird (*P. gigas*), the Yellow-winged

blackbird (*A. thilius*), and the Short-eared owl (*A. flammeus*), show counterintuitive results. They have high indexes of sensitivity to landscape change despite inhabiting open canopy habitats, which is the habitat increased by landclearing. However, these three species have very small clutch sizes and open nests. Further, the giant hummingbird is a reproductive migrant, and the yellow-winged blackbird and the short-eared owl nest on (or very close to) the ground, variables that increase their vulnerability to trampling, parasitism, and predation. Observational and experimental evidence shows that local predators are conspicuous egg consumers (Lazo & Anabalón 1992, Bresciano et al. 1999).

Habitat specialization increases the sensitivity of birds and mammals in mediterranean Chile, which agrees with the reputed most common cause of species endangerment in the region: habitat disturbance. Habitat alteration, largely woodland reduction, accounts for a higher proportion of threatened species than in other parts of the country. Such reduction has been associated to the high intensity of the human occupation of the area (Miller 1980, Simonetti 1999). A reduction in woodlands however, might also increase the area suitable for open canopy dwellers and habitat generalists. Species such as the olivaceous field mouse (*A. olivaceus*) and the degu (*O. degus*) are most common in open shrublands (Glanz 1977, 1984). Their current abundance in central Chile might be a by-product of anthropogenic land use (Simonetti 1989a). None of these species is included in the Red List of Chilean vertebrates, but often they are regarded as pests (Rodríguez 1993).

The sensitivity index, derived independently from any assessment of conservation status, agrees well with the rank of vulnerability derived from the Red List of Chilean vertebrates (see Glade 1993). Although the Red List information does not focus on landscape or habitat modification explicitly, the listing is expected to be indicative of the degree of species sensitivity to any kind of anthropogenic perturbation, including landscape changes. The corre-

spondence between SI and the conservation status strongly suggests that criteria, albeit unstated, used to include species in the Red List are correlated with biological attributes of the species considered. If anything, the Red List fails by being too conservative. While rodent species with high SI were also those species with recognized problems of conservation, several bird species previously considered with no conservation problems according to the Red List (see Glade 1993), had high SI. These avian species included the Giant hummingbird (*P. gigas*), the Yellow-winged blackbird (*A. thilius*), and two rhinocryptids, the White-throated tapaculo (*S. albicolis*), and the Andean tapaculo (*S. magellanicus*). They are all characterized by low reproductive effort, and open nesting behavior (with the exception of *S. albicolis*). Therefore, unless using a more objective, biologically explicit criteria, the Red List can overlook possible conservation problems in species intuitively considered as resistant to disturbance. The sensitivity index can assist specialists in defining species of conservation concern.

The simple sensitivity index used here, after Hansen & Urban (1992), exemplifies that a minimum of biological information about species can provide useful guidelines for the conservation of mediterranean mammals and birds of Chile. Although quantitative genetic studies show that life history traits are more closely linked to fitness than morphological and behavioral traits (Mousseau & Roff 1987, Stearns 1992, Roff 1997), we believe the approach could be extended to include the latter, in particular when there is evidence of conspicuous differences between species, as it is the case of birds (Cody 1974) and rodents (Simonetti 1989b, Vásquez 1996) from central Chile. We support the common plea (e.g., Jaksic & Simonetti 1987) of the need to obtain more detailed information on different aspects of species' life history, read natural history, for accurate determination of the actual and potential effects of landscape transformation on species survival, and hence to protect the biological richness of mediterranean ecosystems.

ACKNOWLEDGMENTS

We thank MTK Arroyo for inviting us to participate in the international workshop "Criterios e Indicadores para la Conservación de la Biodiversidad de los Ecosistemas Mediterráneos en los Países Iberoamericanos", held in Santiago, Chile, during May, 1997. This study was supported by FONDECYT 3950023 (RAV).

LITERATURE CITED

- ARAYA B & G MILLIE (1988) Guía de campo de las aves de Chile, 2nd edition. Editorial Universitaria, Santiago, Chile. 405 pp.
- ARROYO MTK, L CAVIERES, C MARTICORENA & C MUÑOZ (1995) Convergence in the mediterranean floras of central Chile and California: insights from comparative biogeography. In: Arroyo MTK, M Fox & P Zedler (eds) Ecology and biogeography of Mediterranean ecosystems: 43-88. Springer-Verlag, New York.
- BRESCIANO D, JA SIMONETTI, AA GREZ (1999) Edge effects in a mediterranean woodland of central Chile. *Journal of Mediterranean Ecology* 1: 35-40.
- CHARLESWORTH B (1994) Evolution in age-structured populations, 2nd edition. Cambridge University Press, Cambridge. xiv + 306 pp.
- CODY ML (1974) Competition and the structure of bird communities. Princeton University Press, Princeton, New Jersey. x + 318 pp.
- COFRE H & PA MARQUET (1999) Conservation status, rarity, and geographic priorities for conservation of Chilean mammals: an assessment. *Biological Conservation* 88:53-68
- COWLING RM, PW RUNDEL, BB LAMONT, MTK ARROYO & M ARIANOUTSOU (1996) Plant diversity in mediterranean-climate regions. *Trends in Ecology and Evolution* 11: 362-366.
- ESTADES CF (1995) Estimación de la densidad de una comunidad de aves de espinal mediante transectos y estaciones puntuales. *Boletín Chileno de Ornitología* 2: 29-34.
- FAHRIG L & AA GREZ (1996) Population spatial structure, human-caused landscape changes and species survival. *Revista Chilena de Historia Natural* 69: 5-13.
- FALCONER DS & TFC MACKAY (1996) An introduction to quantitative genetics, 4th edition. Longman, London. xvi + 646 pp.
- FORMAN RTT (1995) Land mosaics: the ecology of landscapes and regions. Cambridge University Press, Cambridge. xx + 632 pp.
- FUENTES ER (1990) Landscape change in mediterranean-type ecosystem habitats of Chile: pattern and processes. In: Zonneveld IS & RTT Forman (eds) Changing landscapes: an ecological perspective: 165-190. Springer-Verlag, New York.
- GILPIN ME & ME SOULE (1986) Minimum viable populations: processes of species extinction. In: Soulé ME (ed) Conservation biology: the science of scarcity and diversity: 19-34. Sinauer, Sunderland, Massachusetts.
- GLADE AA, ed (1993) Red list of Chilean terrestrial vertebrates, 2nd edition. Chilean Forest Service, Santiago. viii + 70 pp.
- GLANZ WE (1977) Comparative ecology of small mammal communities in California and Chile. PhD dissertation, University of California, Berkeley, California. 300 pp.
- GLANZ WE (1984) Ecological relationships of two species of *Akodon* in central Chile. *Journal of Mammalogy* 65: 433-441.
- GROOM MJ & N SCHUMAKER (1993) Evaluating landscape change: patterns of worldwide deforestation and local fragmentation. In: Kareiva PM, JG Kinsolver & RB Huey (eds) Biotic interactions and global change: 22-44. Sinauer, Sunderland, Massachusetts.
- GROOMBRIDGE B, ed (1992) Global biodiversity: status of the Earth's living resources. Chapman & Hall, London. xx + 594 pp.
- HANNAH L, JR CARR & A LANKERANI (1995) Human disturbance and natural habitat: a biome level analysis of a global data set. *Biodiversity and Conservation* 4: 128-155.
- HANSEN AJ & DL URBAN (1992) Avian response to landscape pattern: the role of species' life histories. *Landscape Ecology* 7: 163-180.
- JAKSIC FM & JE JIMENEZ (1986) The conservation status of raptors in Chile. *Birds of Prey Bulletin* 3: 95-104.
- JAKSIC FM & JA SIMONETTI (1987) Predator/prey relationships among terrestrial vertebrates: an exhaustive review of studies conducted in southern South America. *Revista Chilena de Historia Natural* 60: 221-244.
- JOHNSON AW (1965) The birds of Chile and adjacent regions of Argentina, Bolivia, and Peru: volume I. Platt Establecimientos Gráficos, Buenos Aires, Argentina. 398 pp.
- JOHNSON AW (1967) The birds of Chile and adjacent regions of Argentina, Bolivia, and Peru: volume II. Platt Establecimientos Gráficos, Buenos Aires, Argentina. 447 pp.
- LAZO I & ANABALON J (1992) Dinámica reproductiva de un conjunto de aves paseriformes de la sabana de espinos de Chile central. *Ornitología Neotropical* 3: 57-64.
- LEEMANS R & G ZUIDEMA (1995) Evaluating changes in land cover and their importance for global change. *Trends in Ecology and Evolution* 10: 76-80.
- MACE GM (1995) Classification of threatened species and its role in conservation planning. In: Lawton JH & RM May (eds) Extinction rates: 198-213. Oxford University Press, Oxford.
- MANN G (1978) Los pequeños mamíferos de Chile: marsupiales, quirópteros, edentados y roedores. *Gayana (Zoología)* 40: 1-342.
- MARTÍNEZ DR & FM JAKSIC (1996) Habitat, relative abundance, and diet of Rufous-legged owls (*Strix rufipes*) King in temperate forest remnants of southern Chile. *Ecoscience* 3: 259-263.
- MCMAMARA JM & AI HOUSTON (1996) State-dependent life histories. *Nature* 380: 215-221.

- MESERVE PL & E LE BOULENGE (1987) Population dynamics and ecology of small mammals in the northern Chilean semiarid region. *Fieldiana Zoology* 39: 413-431.
- MESERVE PL, JR GUTIERREZ, JA YUNGER, LC CONTRERAS & FM JAKSIC (1996) Role of biotic interactions in a small mammal assemblage in semiarid Chile. *Ecology* 77: 133-148.
- MILLER S (1980) Human influences on the distribution and abundance of wild Chilean mammals: prehistoric-present. PhD dissertation, University of Washington, Seattle, Washington. xi + 431 pp.
- MOUSSEAU TA & DA ROFF (1987) Natural selection and the heritability of fitness components. *Heredity* 59: 181-197.
- MUÑOZ A & R MURUA (1990) Control of small mammals in a pine plantation (central Chile) by modification of the habitat of predators (*Tyto alba*, Strigiforme and *Pseudalopex* sp., Canidae). *Acta Oecologica* 11: 251-261.
- MURUA R & J RODRIGUEZ (1989) An integrated control system for rodents in pine plantations in central Chile. *Journal of Applied Ecology* 26: 81-88.
- MYERS N (1990) The biodiversity challenge: expanded hot-spots analysis. *The Environmentalist* 10: 243-256.
- O'CONNOR RJ (1985) Behavioural regulation of bird populations: a review of habitat use in relation to migration and residency. In: Sibly RM & RH Smith (eds) *Behavioural ecology: ecological consequences of adaptive behaviour*: 105-142. Blackwell Scientific Publications, Oxford.
- PRICE T & SCHLUTER D (1991) On the low heritability of life-history traits. *Evolution* 45: 853-861.
- REDFORD KH & JF EISENBERG (1992) *Mammals of the neotropics, the southern cone, volume 2: Chile, Argentina, Uruguay, Paraguay*. The University of Chicago Press, Chicago. x + 430 pp.
- RODRIGUEZ JA (1993) *Roedores plaga: un problema permanente en América Latina y el Caribe*. Oficina Regional de la FAO para América Latina y el Caribe, Santiago. 130 pp.
- ROFF DA (1992) *The evolution of life histories: theory and analysis*. Chapman and Hall, London. xii + 535 pp.
- ROFF DA (1997) *Evolutionary quantitative genetics*. Chapman and Hall, London. xvi + 493 pp.
- SCHLATTER RP (1979) *Avances de la ornitología en Chile*. *Archivos de Biología y Medicina Experimentales* 12: 153-168.
- SIBLY RM & P CALOW (1985) Classification of habitats by selection pressures: a synthesis of life-cycle and *r/K* theory. In: Sibly RM & RH Smith (eds) *Behavioural ecology: ecological consequences of adaptive behaviour*: 75-90. Blackwell Scientific Publications, Oxford.
- SIMONETTI JA (1989a) Small mammals as paleo-environmental indicators: validation for species of central Chile. *Revista Chilena de Historia Natural* 62: 109-114.
- SIMONETTI JA (1989b) Microhabitat use by small mammals in central Chile. *Oikos* 56: 309-318.
- SIMONETTI JA (1999) Diversity and conservation of terrestrial vertebrates of mediterranean Chile. *Revista Chilena de Historia Natural*: this volume.
- SIMONETTI JA & LE CORNEJO (1990) Economic and ecological changes: the prehistory of the Andean mountains of central Chile. In: *Economic catalysts to ecological change*. Working Papers, Center for Latin American Studies, University of Florida, Gainesville: 65-77.
- SIMONETTI JA & B SAAVEDRA (1998) Holocene variation in the small mammal fauna of central Chile. *Zeitschrift für Säugetierkunde* 63: 58-62.
- SOUTHWOOD TRE (1977) Habitat, the templet for ecological strategies? *Journal of Animal Ecology* 46: 337-365.
- SOUTHWOOD TRE (1988) Tactics, strategies and templets. *Oikos* 52: 3-18.
- STEARNS SC (1992) *The evolution of life histories*. Oxford University Press, Oxford. xii + 249 pp.
- SUTHERLAND WJ & PM DOLMAN (1994) Combining behaviour and population dynamics with applications for predicting the consequences of habitat loss. *Proceedings of the Royal Society London Serie B* 255: 133-138.