

# Temporal variability of the sand beach macroinfauna in south-central Chile

Variabilidad temporal de la macroinfauna de playas arenosas en el centro-sur de Chile

E. JARAMILLO<sup>1</sup>, R. STEAD, P. QUIJON, H. CONTRERAS and M. GONZALEZ

Instituto de Zoología, Universidad Austral de Chile, Casilla 567, Valdivia, Chile

<sup>1</sup> E - mail: ejaramil@valdivia.uca.uach.cl

## ABSTRACT

Two sandy beaches were studied in south - central Chile (ca., 39° S), during 17 months to analyze the seasonal variability in population abundances and zonation of the most common species of the intertidal macroinfauna. The beaches displayed differences in mean grain size, Dean's parameter (a composite index of wave and sediment characteristics) and beach face slope. The peracarids *Orchestoidea tuberculata* Nicolet (Amphipoda, Talitridae), *Excirologa braziliensis* Richardson and *Excirologa hirsuticauda* Menzies (Isopoda, Cirolanidae), and the anomuran crab *Emerita analoga* (Stimpson) (Hippidae) were the most common organisms at both beaches, being *E. analoga* and *E. hirsuticauda* the top contributors to abundance. *O. tuberculata* tended to occur in higher abundance, either during autumn - early winter or late summer - early autumn; *E. braziliensis* was more abundant during summer - early autumn, while the abundances of *E. hirsuticauda* and *E. analoga* were quite erratic throughout seasons. In general, the temporal patterns of abundance did not show significant correlations with the beach physical factors considered. The species showed a clear pattern of zonation: *O. tuberculata* occurred in the upper beach levels (around the drift line); *E. braziliensis* occurred in the mid - upper levels of the retention zone; *E. hirsuticauda* occurred throughout all the retention zone and also in the upper and middle levels of the resurgence zone, while *E. analoga* was mostly found in the swash zone and lower levels of the resurgence zone. Although, seasonal changes in the intertidal distribution was observed, it can be seen that each physical zone of the beaches is biologically distinct through seasons.

**Key words:** sandy beaches, macroinfauna, temporal variability, south - central Chile.

## RESUMEN

Se estudiaron dos playas arenosas en el centro sur de Chile (ca., 39° S), con el objetivo de analizar la variabilidad estacional en las abundancias poblacionales y zonación de las especies más comunes de la macroinfauna intermareal. Estas playas mostraron diferencias en tamaño de partícula, parámetro de Dean (índice compuesto de características del oleaje y el sedimento) e inclinación. Los peracáridos *Orchestoidea tuberculata* Nicolet (Amphipoda, Talitridae), *Excirologa braziliensis* Richardson y *Excirologa hirsuticauda* Menzies (Isopoda, Cirolanidae), y el anomuro *Emerita analoga* (Stimpson) (Hippidae) fueron los organismos más comunes en ambas playas, siendo *E. analoga* y *E. hirsuticauda* los mayores contribuyentes a la abundancia. *O. tuberculata* tendió a ocurrir en mayor abundancia, ya sea durante otoño - inicio del invierno o desde fines del verano a inicios del otoño; *E. braziliensis* fue más abundante durante verano - inicios del otoño, mientras que las abundancias de *E. hirsuticauda* y *E. analoga* fueron bastante erráticas a través de las estaciones del año. En general, los patrones temporales de abundancia no mostraron correlaciones significativas con los factores físicos considerados. Las especies mostraron un patrón claro de zonación: *O. tuberculata* ocurrió en los niveles superiores de la playa; *E. braziliensis* ocurrió en los niveles medios y superiores de la zona de retención; *E. hirsuticauda* ocurrió a través de toda la zona de retención y también en los niveles superiores y medios de la zona de resurgencia, mientras que *E. analoga* fue mayoritariamente recolectada en la zona de resaca y niveles inferiores de la zona de resurgencia. Aun cuando se observaron cambios estacionales en la distribución de la macroinfauna, se puede concluir que cada zona física de la playa es biológicamente distinta a través de las estaciones del año.

**Palabras clave:** Playas arenosas, macroinfauna, variabilidad temporal, centro sur de Chile.

## INTRODUCTION

Variation in time and space is a characteristic of ecological systems (Wiens 1986). As stated by Chesson (1986), the common occurrence of this variation and that related to

species populations raises important questions such as How related is the structure of a community to environmental variation? Studies carried out in different habitats have shown that indeed, species and communities are affected by such a variation

(Brewer 1979, Diamond & Case 1986). For example, a decline in biomass of ground finches and seeds was found at an island of the Galápagos during an extended dry period (Grant 1986). Similarly, mean annual precipitation affects seed resources and thus population abundance of desert granivores (Brown et al. 1979). In coastal habitats, it has been found that annual changes in environmental temperature affect the soft bottom macrofauna inhabiting coastal North Sea areas (Beukema 1990). Thus, in the soft bottom bivalves *Cerastoderma*, *Mya* and *Macoma* recruitment on tidal flats of the Wadden Sea is generally poor after mild winters and high after cold winters (Jensen & Jensen 1985, Beukema 1982, 1992). The bivalve *Scrobicularia plana* shows latitudinal variation in reproduction pattern along a coastal gradient extended from the German and Dutch Wadden Sea to France; in this case temperature variation and sediment instability are involved in that population variation (Essink et al. 1991).

Sandy beaches are amongst the most dynamic aquatic habitats (Swart 1983, Brown & McLachlan 1990). Spatial and temporal variations in wave climates (e.g. Aubrey 1983) and seaward gradients in physical conditions (Fleischack & Freitas, 1989, Jaramillo 1987a, McLachlan et al., 1984) confer to these habitats its dynamic characteristics. Thus, it is reasonable to think that in such a dynamic habitat, macrofaunal communities should be influenced by such physical variations. For example, seasonal changes of beach macroinfauna have been related to seasonal changes in beach morphology (e.g., Ansell et al. 1972, McLusky et al. 1975). On the other hand, seasonal variation in macrofauna have been also related to seasonal variation in temperature. For example, Salvat (1966) and Fish (1970) found that in sandy beaches of France and Wales, the abundance of the isopod *Excirrolana pulchra* Leach decreased at the upper levels of its distribution during summer. Jones (1970) observed that in sandy beaches of the south of Wales, the cirrolanid isopods *Eurydice pulchra* Leach and *Eurydice affinis* Hansen, escaped from extreme winter temperatures by moving from intertidal to subtidal levels. Seasonal variation in

temperature appears to affect decapod movements, and consequently seasonal variation in community structure in a sandy beach of North Carolina (Leber 1982).

Studies concerning the Chilean sandy beach macroinfauna have been mainly devoted to describe zonation patterns of the intertidal macroinfauna. The first one was that of Dahl (1952) who described the across shore distribution of crustaceans at Montemar, Puerto Montt and Punta Arenas (central, south central and southern extreme of Chile, respectively). Afterwards, the sandy beach macroinfauna of central Chile was studied by Osorio et al. (1967) and Nuñez et al. (1974), that of the north by Jaramillo (1987b) and Clarke & Peña (1988), that of the north central Chile by Sanchez et al. (1982) and Jaramillo (1987 b), and that of south central Chile by Jaramillo (1978, 1982, 1987 a, b, 1994), Jaramillo & González (1991), Jaramillo & McLachlan (1993), and Jaramillo et al. (1993). The vast majority of these studies are the result of short - term surveys. Even though, two general conclusions can be drawn from these studies: i) the numerically dominant organisms along the Chilean coast are crustaceans (mostly Peracarida and Anomura), and ii) peracarid species typically dominate upper and middle beach levels, while anomurans dominate at the lower beach levels.

Jaramillo & McLachlan (1993), Jaramillo et al. (1993) and Jaramillo (1994) have shown that community parameters (species richness, abundance and biomass) and zonation of the sandy beach macroinfauna of south central Chile, are closely related to the morphodynamic type of the beach measured by Dean's parameter (Short & Wright 1983). However, the studies reported so far for this coast have not addressed the effect of time (e.g. seasonal changes) on beach morphodynamics, and consequently upon community parameters and zonation of the macroinfauna.

Sandy beaches of south central Chile are characterized by seasonal changes in beach topography and grain size (Jaramillo 1987 a). Winter throughout spring is typically an erosion period, while summer and autumn are characterized by accretion of sands which becomes finer than during the former period

(Jaramillo 1987a). Given the temporal pattern of sand variability we hypothesized that macroinfaunal temporal changes at two sandy beaches located in south central Chile (Mehuín, ca. 39° S) follow beach physical changes.

#### MATERIAL AND METHODS

The two beaches studied were located at Mehuín, in the south central area of the Chilean coast (approximately 39 - 40° S, Fig. 1). These beaches are separated by a rocky shore promontory; Playa Grande is a beach 2000 m in length, while the other one (Playa Universitaria) is a beach approximately 300 m long (Fig. 1). We chose two closely

located beaches to avoid eventual effects of regional variability on the conclusions of the study.

The morphodynamic type of both beaches correspond to the high-energy intermediate to dissipative category of Short & Wright (1983). They are fully exposed to the breaking waves of the Pacific Ocean. The tides are semidiurnal with a maximum range of approximately 1.5 m.

Study sites were sampled monthly during spring low tides from April 1992 to August 1993. On each site, samples (0.03 m<sup>2</sup>, 30 cm deep) were collected with plastic cores at ten equally spaced levels along three replicated transects (1 m apart each other) extending from above the drift line to the swash zone; i.e., the uppermost station was located above




Fig. 1: Location of the sandy beaches studied at Mehuín, south - central Chile: Playa Universitaria (PU) and Playa Grande (PG).

Localización de las playas arenosas estudiadas en Mehuín, centro-sur de Chile: Playa Universitaria (PU) y Playa Grande (PG).

the drift line, the second on the drift line and the last at the lowest limit of the swash zone (indicated by bore collapse). The sediment was sieved through 1 mm mesh and the macroinfauna was stored in 5% formalin until sorting. Abundance values per running meter of beach were obtained by linear interpolation between sampling stations, after obtaining mean abundances per m<sup>2</sup> at each station. The temporal variation in the zonation of the macroinfauna was analyzed by visual examination of kite diagrams and temporal variation of distribution modes (i.e. location of the population mode).

Wave height was estimated on every visit by measuring the height of breaking waves ( $n = 10$ ) with graduated poles against the horizon, and adding the result to the height difference between the location of the observer and the lowest point where the backwash met the next incoming swash bore. The wave period (measured with a stop watch,  $n = 10$ ) was measured as the time interval between breakers. Triplicate sediment samples (1 m apart each other) were taken from the upper, middle and lower beach (stations 2, 5 and 10, respectively), for grain size analysis; which was carried out using a settling tube (Emery 1938). Mean grain size was calculated according to the moments computational method (Seward - Thompson & Hails 1973) and used to estimate sand fall velocity according to the method described by Gibbs et al. (1971). From estimated mean wave height, wave period and sand fall velocity, Dean's dimensionless parameter ( $\Omega$ ) (Short & Wright 1983) was calculated for each sampling day:  $\Omega = \text{wave height (cm)} / \text{sand fall velocity (cm s}^{-1}\text{)} \times \text{wave period (s)}$ . The morphology (i.e., beach face slope at the site of the transect) of each site was determined by Emery's profiling technique (Emery 1961).

Sediment samples were collected at each station for water content analyses by using a 3.5 cm diameter plastic core (3 cm of depth). Samples were wrapped in aluminium foil and weighed immediately after collection. Water content of the sand was estimated as the loss in weight of wet sediments after drying (120°C for 96 hours). The penetrability of the sediments was measured by dropping a 33.6 g metal rod down a 1 m tube. The depth to

which the rod penetrated into the sediment was measured 4 times at each station.

Visual observations of the variability in moisture across the intertidal were used to analyze variability in zonation of the most common species. The visual observations were carried out to observe the limits of the dry, retention, resurgence, and swash (= saturation) zones, as defined by Salvat (1964, 1966). In this categorization the dry zone corresponds to that located at the top of the shore (usually above the drift line) where the sand is normally wetted only by spray; the retention zone is that area of the intertidal which is reached by all tides and where some moisture always remains in the interstitial space; the resurgence zone is the zone marked by movement of water on the outgoing tide, and the saturation zone is that permanently saturated with water (McLachlan 1990, Salvat 1964, 1966)

## RESULTS

### *The beaches*

Figure 2 and Table 1 show the variability in physical factors at the two beaches. Wave tended to be higher during autumn and winter and lower during summer. No significant differences between beaches were found when the averages based on 17 monthly data of wave height were compared (Table 1). The same was found for wave periods (Table 1). In general, the coarsest sands were found at Playa Universitaria (Fig. 2); indeed, the average for this parameter (274.1  $\mu\text{m}$ ) at this beach was significantly higher (Table 1).

The temporal pattern of Dean's parameter ( $\Omega$ ) at both beaches showed similar trends. During some months, both beaches had  $\Omega$  values lower than 6 which is indicative of intermediate beach types (*sensu* Short & Wright 1983); on the other hand, during other months,  $\Omega$  values were higher ( $> 8$ ) which is typical of beaches displaying dissipative characteristics (*sensu* Short & Wright 1983). Generally, the highest Dean's values were estimated when wave heights were higher. Taking into account the average based on 17 monthly data it was found that the Dean's parameter was significantly




Fig. 2: Temporal variability in wave (breaker) height (Hb), wave period (T), mean particle size ( $\mu\text{m}$ ), Dean's parameter ( $\Omega$ ) and beach face slope ( $1/\text{slope}$ ).

Variabilidad temporal de altura de las olas en zona de rompiente (Hb), período de la ola (T), tamaño medio de la partícula ( $\mu\text{m}$ ), parámetro de Dean ( $\Omega$ ) y pendiente de la cara de la playa ( $1/\text{slope}$ ).

higher at Playa Grande (i.e., a more dissipative condition) (Table 1). This beach also had flatter slopes, especially during June - July 1992 (Fig. 2).

#### Abundance of the macroinfauna

As found in other studies (Jaramillo & McLachlan 1993, Jaramillo et al. 1993, Jaramillo 1994), the peracarids *Orchestoidea tuberculata* Nicolet (Amphipoda, Talitridae), *Exciorolana braziliensis* Richardson and *Exciorolana hirsuticauda* Menzies (Isopoda, Cirolanidae), and the anomuran crab *Emerita*

*analoga* (Stimpson) (Hippidae) were the most common organisms at both beaches, being *Emerita analoga* and *Exciorolana hirsuticauda* the most abundant (Fig. 3).

The temporal variability in population abundances of these species is shown in Figure 3. *Orchestoidea tuberculata* tended to occur in higher abundance, during autumn - early winter and late summer - early autumn; this pattern is more evident at Playa Universitaria where the highest abundance occurred during April - July 1992 and March - May 1993. *Exciorolana braziliensis* was more abundant during summer - early autumn (February - April); while the abundance of *Exciorolana hirsuticauda* and *Emerita analoga* were the most erratic, with many peaks being observed throughout the study period (Fig. 3).

The temporal variability in the abundance of *Exciorolana braziliensis* was positively correlated with Dean's parameter (i.e., the highest abundances of this isopod occurred when Dean's were lower; a more intermediate beach condition) and negatively correlated with beach face slopes. By contrast, other species did not show any significant correlations (Table 2).

The abundance of *Orchestoidea tuberculata* was significantly higher at Playa Universitaria during the first four months of the study period, but there were not differences in the rest of the period. In general, the other species did not show differences between the two beaches (Fig. 3).

#### Zonation of the macroinfauna

The monthly intertidal distribution of *Orchestoidea tuberculata*, *Exciorolana braziliensis*, *Exciorolana hirsuticauda* and *Emerita analoga* is shown in Figs. 4-7. Figure 8 shows the monthly variation in the location of the distribution mode of these species.

The amphipod *Orchestoidea tuberculata* occurred higher up on the beach (Fig. 4). The general trend was to show its maximum abundance above the drift line (dry zone) during late autumn throughout early spring (May - October), and below it (i.e., at the upper levels of the retention zone) during the summer months (December - March; Fig. 4). The analysis of the monthly variability of the

TABLE I

Physical characteristics at each beach and results of one way analysis of variance. Averages based on monthly data (n = 17 months). See values of p in the results of variance analyses  
 Características físicas de cada playa y resultados de los análisis de varianza de una vía. Promedios basados en datos mensuales (n = 17 meses). Ver valores de p en los resultados de los análisis de varianza

ANALYSIS OF VARIANCE								
Variables	Beaches	Average	Source of variation	ss	df	ms	f ratio	p
wave height (cm)	Playa Universitaria	297.6	between groups	10658.9	1	10658.9	0.947	0.3480
	Playa Grande	333.0	within groups	360022.1	32	11250.7		
			total	370681.1	33			
wave period (sec.)	Playa Universitaria	14.6	between groups	11.1	1	11.1	0.915	0.3561
	Playa Grande	13.4	within groups	387.1	32	12.1		
			total	398.2	33			
mean grain size ( $\mu\text{m}$ )	Playa Universitaria	274.1	between groups	6062.2	1	6062.2	8.864	0.0055
	Playa Grande	247.4	within groups	21884.8	32	683.9		
			total	27947.1	33			
$\Omega$	Playa Universitaria	5.7	between groups	43.1	1	43.1	5.926	0.0207
	Playa Grande	8.0	within groups	233.0	32	7.3		
			total	276.1	33			
1/slope	Playa Universitaria	26.9	between groups	2439.5	1	2439.5	7.653	0.0093
	Playa Grande	43.9	within groups	10200.7	32	318.8		
			total	12640.2	33			




Fig. 3: Temporal variability in macroinfaunal abundances.

Variabilidad temporal en la abundancia de la macroinfauna.

distribution mode shows that at Playa Universitaria, *Orchestoidea tuberculata* moved up to beach levels located above the drift line during autumn and winter months and to lower beach levels (close to the drift line) during the warmer months (Fig. 8). A similar pattern of seasonal movements was not observed at Playa Grande where the distribution mode was always near by to the drift line (Fig. 8).

The isopod *Excirolana braziliensis* occurred in the mid and upper levels of the retention zone (right below the drift line) without showing much seasonal variability in its zonation (Fig. 5). *Excirolana hirsuticauda* occurred throughout the entire retention zone; during some months, this isopod was also found in the upper and middle levels of the resurgence zone (Fig. 6). The distribution modes of both isopods was quite shifting at both beaches; it were usually found below the drift line (Fig. 8).

The anomuran crab *Emerita analoga* was mostly found in the swash zone and lower levels of the resurgence zone (Fig. 7). At both beaches, and during winter and spring, the distribution mode of *Emerita analoga* shifted towards lower beach levels than

TABLE 2

Results of regression analyses for the abundances of the species versus  $\Omega$  and beach face slope from each site

Resultados de los análisis de regresión para las abundancias de las especies versus  $\Omega$  y pendiente de cada sitio

Beaches	Species	$\Omega$				1/slope			
		a	b	r	p	a	b	r	p
Playa Universitaria	<i>Orchestoidea tuberculata</i>	7018.35	112.80	0.05	0.82	-2844.66	390.06	0.65	0.01
	<i>Excirolana braziliensis</i>	6761.26	-664.19	-0.48	0.05	-2844.09	202.36	0.47	0.05
	<i>Excirolana hirsuticauda</i>	5043.70	354.28	0.22	0.40	5905.23	43.29	0.08	0.74
	<i>Emerita analoga</i>	30324.90	-847.96	-0.12	0.64	8082.48	645.45	0.30	0.24
Playa Grande	<i>Orchestoidea tuberculata</i>	4265.03	-141.21	-0.15	0.56	3722.65	-13.49	-0.12	0.66
	<i>Excirolana braziliensis</i>	2545.88	211.01	0.24	0.35	3907.97	7.32	0.07	0.80
	<i>Excirolana hirsuticauda</i>	15914.40	-224.13	0.08	0.76	8314.31	132.45	0.38	0.13
	<i>Emerita analoga</i>	18175.90	513.43	0.12	0.63	10558.80	266.91	0.52	0.03




Fig. 4: Intertidal distribution of the amphipod *Orchestoidea tuberculata* relative to the drift line (DL). Numbers in the y axis refer to cm above and below drift line.

Distribución intermareal del anfípodo *Orchestoidea tuberculata* en relación a la línea de resaca. Los números en el eje y se refieren a cm sobre y bajo la línea de resaca.

those occupied during some warmer months (Fig. 8).

*Orchestoidea tuberculata* and *Excirolana braziliensis* occurred in sediments which rendered similar ranges in water content at both beaches (ca. 1 - 21%) (Fig. 9, Table 3). However, the most frequent occurrences for the former species were in sands with higher water contents (see the mode for both species at both beaches in Table 3). *Excirolana hirsuticauda* was collected from sediments with water contents between approximately 4

and 23% (Fig. 9); most frequently it occurred in sediments having 16 - 19% of water content (see the modes in Table 3). *Emerita analoga* was primarily collected from sediments having 20 - 21 % of water content (Table 3); its range was 8 - 34% (Fig. 9).

The results presented in Fig. 10 and Table 3 show that the sediments occupied by *Orchestoidea tuberculata* and *Excirolana braziliensis* were the softer ones (i.e., more penetrable during the low tide samplings), while those in which *Excirolana hirsuticauda*




Fig. 5: Intertidal distribution of the isopod *Exciorolana braziliensis* relative to the drift line (DL). Numbers in the y axis refer to cm above and below drift line.

Distribución intermareal del isópodo *Exciorolana braziliensis* en relación a la línea de resaca. Los números en el eje y se refieren a cm sobre y bajo la línea de resaca.

TABLE 3

Water content and penetrability values of sediments from which the species were collected, in the beaches Playa Universitaria and Playa Grande

Contenidos de agua y valores de penetrabilidad en los sedimentos desde donde se recolectaron las especies en Playa Universitaria y Playa Grande

Beaches	Species	Water content (%)				Penetrability (cm)			
		Average	(s.d.)	Mode	Range	Average	(s.d.)	Mode	Range
Playa Universitaria	<i>Orchestoidea tuberculata</i>	6.8	(4.7)	5.1	1.3-20.1	2.7	(1.1)	2.1	0.9-6.0
	<i>Exciorolana braziliensis</i>	7.3	(5.4)	4.7	1.3-19.2	2.6	(1.2)	2.1	1.0-6.0
	<i>Exciorolana hirsuticauda</i>	18.0	(3.8)	16.0	3.9-23.0	1.2	(0.1)	1.0	0.6-2.7
	<i>Emerita analoga</i>	21.5	(3.1)	21.0	8.4-33.3	1.6	(0.5)	1.2	0.8-3.1
Playa Grande	<i>Orchestoidea tuberculata</i>	11.9	(5.4)	5.5	1.3-21.2	2.0	(0.9)	1.8	0.4-4.8
	<i>Exciorolana braziliensis</i>	12.5	(6.0)	4.3	1.3-21.3	1.9	(0.9)	1.6	0.7-4.8
	<i>Exciorolana hirsuticauda</i>	17.4	(3.9)	19.3	3.8-22.7	1.1	(0.4)	0.9	0.6-3.8
	<i>Emerita analoga</i>	21.9	(3.0)	20.2	13.3-33.8	1.3	(0.4)	1.2	0.6-2.5

and *Emerita analoga* were collected were the harder ones (i.e., lower penetrability values).

#### DISCUSSION

This study has shown that the physical characteristics of the beaches studied are quite dynamics. Consequently, it was expected that changes in beach morpho-

dynamics might well affect the macroinfauna. However, habitat variation and macroinfaunal variations were not closely associated, at least on the scale of month to month changes. This suggests that apart from the beach physical factors studied, another ones should be involved in such a temporal variation of the macroinfauna. It is also possible that the macroinfauna of the sandy beaches studied indeed changes rapidly to




Fig. 6: Intertidal distribution of the isopod *Excirolana hirsuticauda* relative to the drift line (DL). Numbers in the y axis refer to cm above and below drift line.

Distribución intermareal del isópodo *Excirolana hirsuticauda* en relación a la línea de resaca. Los números en el eje y se refieren a cm sobre y bajo la línea de resaca.




Fig. 7: Intertidal distribution of the anomuran *Emerita analoga* relative to the drift line (DL). Numbers in the y axis refer to cm above and below drift line.

Distribución intermareal del anomuro *Emerita analoga* en relación a la línea de resaca. Los números en el eje y se refieren a cm sobre y bajo la línea de resaca.

track environmental variations, but those changes occur in a time interval to which our sampling program was not the adequate to find relationships. It should be also pointed

out that the beach state index (Dean's parameter) is strongly affected by wave height, a physical parameter which was estimated during a single observation period




Fig. 8: Temporal variation of the distribution modes of the crustaceans relative to the drift line (DL) at Playa Universitaria (PU) and Playa Grande (PG). Numbers in the y axis refer to cm above and below drift line.

Variabilidad temporal en las modas distribucionales de los crustáceos en relación a la línea de resaca. Los números en el eje y se refieren a cm sobre y bajo la línea de resaca.

at each site. Thus it is possible that this highly temporally variable parameter may have had a strong influence on monthly values calculated for this index, thus rendering misleading results.

The absence of relationships between habitat and faunal variations at the beaches studied means that the relationship found by several authors between beach type and community structure of the macroinfauna (McLachlan 1990, McLachlan et al. 1993, Jaramillo & McLachlan 1993, Jaramillo et al. 1993 and Jaramillo 1994) does not hold when time scales longer than snapshot samplings are considered. However, there are some studies which suggest that there are close relationships between physical and

macroinfaunal variation. For example, Croker et al. (1975) and Leber (1982) reported seasonal changes in the community structure of sandy beaches of northern New England and North Carolina, respectively; changes probably related to seasonal changes in environmental temperature. Brazeiro & Defeo (1996) reached similar conclusions after studying a sandy beach of Uruguay for a period of one year. For seasonal differences in abundances and zonation of the macroinfauna inhabiting a sandy beach of north central Chile, Sánchez et al. (1982) suggested variation in grain size and beach face slope.

The findings of this study show that the zonation patterns described earlier for the macroinfauna of this area (Jaramillo &




Fig. 9: Frequency of water contents of sediments from which *Orchestoidea tuberculata*, *Excirolana braziliensis*, *Excirolana hirsuticauda* and *Emerita analoga* were collected.

Frecuencias de contenidos de agua en los sedimentos de los cuales se recolectaron *Orchestoidea tuberculata*, *Excirolana braziliensis*, *Excirolana hirsuticauda* y *Emerita analoga*.




Fig. 10: Frequency of penetrabilities of sediments from which *Orchestoidea tuberculata*, *Excirolana braziliensis*, *Excirolana hirsuticauda* and *Emerita analoga* were collected.

Frecuencias de penetrabilidades en los sedimentos de los cuales se recolectaron *Orchestoidea tuberculata*, *Excirolana braziliensis*, *Excirolana hirsuticauda* y *Emerita analoga*.

McLachlan 1993, Jaramillo et al. 1993, Jaramillo 1994), and which resulted from snapshot samplings hold quite well. Thus, even when species shift seasonally in their across shore distribution, they were distributed in a well defined zonation pattern with talitrid amphipods and cirrolanid isopods in the upper beach levels, cirrolanid isopods in the middle ones and anomuran crabs in the lower beach levels. It is worthwhile mentioning that this zonation pattern represent the day low tide zonation when most of the animals are buried below the sand surface. This accentuates the physical differences of sand types occupied by the different species and exemplified by the specific differences in water content and penetrability of sands.

The former results are also reminiscent of those reported in the supra regional study of Dahl (1952), the paradigmatic study to which most zonation schemes are compared; the amphipod as a characteristic species of the supratidal zone and the isopods as typical from the midlittoral zone. Dahl (1952) also found anomuran crabs (*Emerita analoga*) in the lower beach levels of a sandy beach located in central Chile (Montemar); however, he did not use this organism to characterize the infralittoral zone of temperate, but of tropical and subtropical beaches.

The results of this study show that the suggested zonation scheme of Salvat (1964, 1966), which is based upon relative humidity of sands, and thus independent of zoogeographical limitations and presence of characteristics organisms, can be superimposed to that of Dahl (1952). *Orchestoidea tuberculata* and *Excirolana braziliensis* occurred highest on the beach, with a trend for the former species to shift a bit higher during some months, thus characterizing the dry upper beach zone with the driest and softest sands. The latter species characterized the upper levels of the retention zone. *Excirolana hirsuticauda* occupied the mid beach levels, primarily the lower levels of the retention zone, while *Emerita analoga* was a typical organism of the lower shore, characterizing the resurgence - swash zone.

But even when both, Dahl's (1952) and Salvat's (1964) schemes can be superimposed themselves for the case of the studied beaches, some exceptions are still found; e.g, in some months the distribution of the isopod *Excirolana braziliensis* extended up to the dry zone - supralittoral zone.

In conclusion, our results show that despite the dynamic nature of the beaches studied, the population abundances of their crustaceans tend to be independent of beach stage. They also tend to have temporal changes, but no always seasonal repeatable distributional patterns. The species studied here, were always present at both study sites. Both, a cirrolanid isopod (*Excirolana hirsuticauda*) and an anomuran crab (*Emerita analoga*) are the most abundant species, a conclusion which seems to stand for all the south central Chile, providing that similar types of

beaches are compared (cf. Jaramillo *et al.*, 1993).

#### ACKNOWLEDGMENTS

We thank Jacqueline Muñoz, Claudia Añazco, Victor Poblete, Robert Brümmer and María Avellanal for assistance with field work. We also thank A. De Ruyck, A. Soares (University of Port Elizabeth, South Africa) and anonymous reviewers for critics and suggestions to an earlier manuscript. This study was made possible by financial support from CONICYT (Chile) (FONDÉCYT research project 92-191) and Universidad Austral de Chile (DID projects S92/36 and S94/30).

#### LITERATURE CITED

- ANSELL AD, DS MCLUSKY, A STIRLING & A TREVALLION (1972). The ecology of two sandy beaches in South West India. I. Seasonal changes in physical and chemical factors and in the macroinfauna. *Marine Biology* 17: 38-62.
- AUBREY DG (1983). Beach changes on coasts with different wave climate. *In: McLachlan, A. & Th. Erasmus (Eds.) Sandy beaches as ecosystems*. W. Junk, The Hague, pp. 63-85.
- BEUKEMA JJ (1982). Annual variations in reproductive success and biomass of the major macrozoobenthic species living in a tidal flat area of the Wadden Sea. *Netherlands Journal of Sea Research* 16: 37-45.
- BEUKEMA JJ (1990). Expected effects of changes in winter temperatures on benthic animals living in soft sediments in coastal North Sea areas. *In: Beukema JJ et al. (Eds.) Expected Effects of Climatic Change on Marine Coastal Ecosystems*. Kluwer Academic Publishers The Netherlands, pp. 83-92.
- BEUKEMA JJ (1992). Expected changes in the Wadden Sea benthos in a warmer world: lessons from period with mild winters. *Netherlands Journal of Sea Research* 30: 73-79.
- BRAZEIRO A & O DEFEO (1996). Macroinfauna zonation in microtidal sandy beaches: is it possible to identify patterns in such variable environments?. *Estuarine, Coastal and Shelf Science* 42: 523-536.
- BREWER R (1979). *Principles of ecology*. Saunders College, Philadelphia, 299 pp.
- BROWN AC & A McLACHLAN (1990). *Ecology of sandy shores*. Elsevier Science Publishers B.V. Amsterdam, 328 pp.
- BROWN JH, OJ REICHMAN & DW DAVIDSON (1979). Granivory in desert ecosystems. *Annual Review of Ecology and Systematics*, 10: 201-227.
- CHESSON PL (1986). Environmental variation and the coexistence of species. *In: Diamond J & TJ Case (Eds.) Community ecology*. Harper & Row, Publishers, New York, 240-256.
- CLARKE M & R PEÑA (1988). Zonación de la macroinfauna en una playa de arena del norte de Chile. *Estudios Oceanológicos* 7: 17-31.
- CROKER RA, RP HAGER & KJ SCOTT (1975). Macroinfauna of northern New England marine sand. II. Amphipod dominated intertidal communities. *Canadian Journal of Zoology*, 53: 42-51.
- DAHL E (1952). Some aspects of the ecology and zonation of the fauna on sandy beaches. *Oikos* 4: 1-27.
- DIAMOND J & TJ CASE (1986). *Community ecology*. Harper & Row, Publishers, New York, 665 pp.
- EMERY KO (1938) Rapid method of mechanical analysis of sands. *Journal of Sedimentary Petrology* 8: 105-111.
- EMERY KO (1961) A simple method of measuring beach profiles. *Limnology and Oceanography* 6: 90-93.
- ESSINK K, JJ BEUKEMA, J COOSEN, JA CRAEYMEERSCH, J-P DUCROTOY, H. MICHAELIS & B ROBINEAU (1991) Population dynamics of the bivalve mollusc *Scrobicularia plana* da Costa: comparisons in time and space. *In: Elliot M & J-P Ducrottoy (eds) Estuaries and coasts: spatial and temporal intercomparisons:167-172*. ECSA 19 Symposium, Olsen & Olsen.
- FISH S (1970) The biology of *Euridice pulchra* (Crustacea: Isopoda). *Journal of Marine Biology Association, U.K.* 50: 753-785.
- FLEISCHACK PC & AJ DE FREITAS (1989) Physical parameters influencing the zonation of surf zone benthos. *Estuarine, Coastal and Shelf Science* 28: 517-530.
- GIBBS RJ, MD MATTHEWS & DA LINK (1971) The relationship between sphere size and settling velocity. *Journal of Sedimentary Petrology* 41: 7-18.
- GRANT PR (1986). Interspecific competition in fluctuating environments. *In: Diamond J & TJ Case (eds) Community ecology*.: 173-191. Harper & Row, Publishers, New York.
- JARAMILLO E (1978) Zonación y estructura de la comunidad macrofaunística en playas de arena del sur de Chile (Mehuín, Valdivia). *Studies on Neotropical Fauna and Environment* 13: 71-92.
- JARAMILLO E (1982) Taxonomy, natural history and zoogeography of sand beach isopods from the coast of southern Chile. *Studies on Neotropical Fauna and Environment* 17: 175-194.
- JARAMILLO E (1987 a) Community ecology of Chilean sandy beaches. Ph.D. Thesis. University of New Hampshire, Durham, U.S.A.
- JARAMILLO E (1987 b) Sandy beach macroinfauna from the Chilean coast: zonation patterns and zoogeography. *Vie et Milieu* 37: 165-174.
- JARAMILLO E & M GONZALEZ (1991) Community structure and zonation of the macroinfauna along a dissipative reflective range of beach category in southern Chile. *Studies on Neotropical Fauna and Environment* 26: 193-212.
- JARAMILLO E & A MCLACHLAN (1993) Community and population responses of the macroinfauna to physical factors over a range of exposed sandy beaches in south-central Chile. *Estuarine, Coastal and Shelf Science* 37: 615-624.
- JARAMILLO E, A MCLACHLAN & P COETZEE (1993) Intertidal zonation patterns of macroinfauna over a range of exposed sandy beaches in south-central Chile. *Marine Ecology Progress Series* 101: 105-118.
- JARAMILLO E (1994) Patterns of species richness in sandy beaches of South America. *South African Journal of Zoology* 29: 227-234.
- JENSEN KT & JN JENSEN (1985). The importance of some epibenthic predators on the density of juvenile benthic macrofauna in the Danish Wadden Sea. *Journal of Experimental Marine Biology and Ecology* 89: 157 - 174.

- JONES DA (1970) Factors affecting the distribution of the intertidal isopods *Eurydice pulchra* Leach and *E. affinis* Hansen in Britain. *Journal of Animal Ecology* 39: 455-472.
- LEBER KM (1982) Seasonality of macroinvertebrates on temperate, high wave energy sandy beach. *Bulletin of Marine Science* 32: 86 - 98.
- McLACHLAN A (1990) Dissipative beaches and macrofauna communities on exposed intertidal sands. *Journal of Coastal Research* 6: 57-71.
- McLACHLAN A, AC COCKCROFT & DE MALAN (1984) Benthic faunal response to a high energy gradient. *Marine Ecology Progress Series* 16: 51-53.
- McLACHLAN A, E JARAMILLO, TE DONN & F WESSELS (1993) Sandy beach macrofauna communities and their control by the physical environment: a geographical comparison. *Journal of Coastal Research (Special Issue)* 15: 27-38.
- McLUSKY DS, SA NAIR, A STIRLING & R NHARGAVA (1975) The ecology of a central west Indian beach, with particular reference to *Donax incarnatus*. *Marine Biology* 30: 267-276.
- NUÑEZ T, O ARACENA & MT LOPEZ (1974) *Emerita analoga* en Llico, Provincia de Curico (Crustacea, Decapoda, Hippidae). *Boletín de la Sociedad de Biología de Concepción (Chile)* 48: 11-22.
- OSORIO C, N BAHAMONDE & MT LOPEZ (1967) El limanche *Emerita analoga* (Stimpson) en Chile. *Boletín del Museo Nacional Historia Natural de Chile* 29: 61-116.
- SALVAT B (1964) Les conditions hydrodynamiques interstitielles des sédiments meubles intertidaux et la repartition verticale de la faune endogée. *Comptes Rendus Academie Sciences Paris* 256: 1576-1579.
- SALVAT B (1966) *Eurydice pulchra* (Leach 1815), et *Eurydice affinis* (Hansen 1905), (Isopodes: Cirolanidae) - taxonomie, éthologie, écologie, repartition verticale et cycle reproducteur. *Actes de la Societe Linneenne de Bordeaux, Serie A N° 1*, 103: 1-77.
- SANCHEZ M, JC CASTILLA & O MENA (1982) Variaciones verano - invierno de la macrofauna de arena en Playa Morrillos (Norte Chico, Chile). *Studies on Neotropical Fauna and Environment* 17: 31-49.
- SEWARD-THOMPSON B & J HAILS (1973) An appraisal on the computation of statistical parameters in grain size analysis. *Sedimentology* 11: 83-98.
- SHORT AD & LD WRIGHT (1983) Physical variability of sandy beaches. In: McLachlan A & Th Erasmus (eds) *Sandy beaches as ecosystems*: 133-144. W. Junk, The Hague.
- SWART DH (1983) Physical aspects of sandy beaches- a review. In: McLachlan A & Th Erasmus (eds) *Sandy beaches as ecosystems*: 5-44. W. Junk, The Hague.
- WIENS JA (1986) Spatial scale and temporal variation in studies of shrubsteppe birds. In: Diamond J & TJ Case (eds) *Community ecology*: 154-172. Harper & Row, Publishers, New York.